



Habitat Type

- Species Present
- Rearing Area
- Spawning Area

Drainage Basin

- Allen Creek
- Ebey Slough
- West Fork Quilceda Creek
- Edgecomb/Hayho Creeks
- Middle Fork Quilceda Creek
- Quilceda Creek
- King Creek

Data: Washington SWIFD 2023



COHO MALE SPAWNING



COHO FEMALE ADULT

TIMING OF COHO FRESHWATER LIFE STAGES

Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Spawning												
Incubation												
Fry												
Juveniles												

Coho salmon utilize shallow reaches and small streams for spawning. The young spend one to two years in fresh water before migration.



THE CITY OF MARYSVILLE DISCLAIMS ANY WARRANTY OF MERCHANTABILITY OR WARRANTY OF FITNESS OF THIS MAP FOR ANY PARTICULAR PURPOSE, EITHER EXPRESSED OR IMPLIED. NO REPRESENTATION OR WARRANTY IS MADE CONCERNING THE ACCURACY, CURRENCY, COMPLETENESS OR QUALITY OF DATA DEPICTED. ANY USER OF THIS MAP ASSUMES ALL RESPONSIBILITY FOR USE THEREOF, AND FURTHER AGREES TO HOLD THE CITY OF MARYSVILLE HARMLESS FROM AND AGAINST ANY DAMAGE, LOSS, OR LIABILITY ARISING FROM ANY USE OF THIS MAP.